

Sea-level databases

Nicole S. Khan¹, F. Hibbert² and A. Rovere³

The study of past sea levels relies on the availability of standardized sea-level reconstructions, which allow for broad comparison of records from disparate locations to unravel spatial patterns and rates of sea-level change at different timescales. Subsequently, hypotheses about their driving mechanisms can be formulated and tested.

Approach to database compilation

Geological sea-level reconstructions are developed using sea-level proxies, which formed in relation to the past position of sea level and include isotopic, sedimentary, geomorphic, archaeological, and fixed biological indicators, in addition to coral reefs and microatolls, as well as wetland flora and fauna. The past position of sea level over space and time is defined by what are termed sea-level index or limiting points, which are characterized by the following fundamental fields: a) geographic location; b) age of formation, traditionally determined by radiometric methods (e.g. radiocarbon or U-series dating); c) the elevation of the sample with respect to a contemporary tidal datum; and d) the relationship of the proxy to sea level at the time of formation (i.e. the proxy's "indicative meaning", which describes the central tendency (reference water level) and vertical (indicative) range) relative to tidal levels. Although conceptually only four primary fields are necessary to define a sea-level index point, in practice many more fields are required to appropriately archive information related to geological samples (e.g. stratigraphic context, sample collection,

laboratory processing), and it is important to distinguish between primary observations and secondary interpretation so that the latter may be updated as science advances (see Hibbert et al. 2016; Hijma et al. 2015).

While this approach was developed through the International Geoscience Programme projects running from the 1970s to present and has been widely applied to Holocene reconstructions (e.g. Shennan and Horton 2002), it has only more recently been adopted for older archives and time periods (e.g. Rovere et al. 2014, 2016). The standardization of sea-level databases of various ages has been one of the main objectives of the PAGES PALEO constraints on SEA level (PALSEA) working group (Düsterhus et al. 2016) and by projects related to it (e.g. the International Union for Quaternary Research (INQUA) Geographic variability of HOLOCENE relative SEA level (HOLSEA) and MEDITERRANEAN sea-level change and projection for future FLOODING (MEDFLOOD) projects). Here we describe recent progress and advances in database compilation, and highlight remaining challenges and future directions.

Last Glacial Maximum to present

The standardization of sea-level databases spanning time periods from the Last Glacial Maximum (LGM) to present has seen rapid development in recent years. Notable progress has been made through a community effort, unified under the HOLSEA project, to develop a standardized global database of post-LGM sea levels. The first iteration of this database was made available in April 2019 through a special issue entitled "Inception of a Global Atlas of Sea Levels since the Last Glacial Maximum" published in *Quaternary Science Reviews*. Regional contributions in the special issue from Atlantic Canada, the British Isles, the Netherlands, Atlantic Europe, the western Mediterranean, Israel, the Russian Arctic, South Africa, the Malaysian Peninsula, and Southeast Asia, India, Sri Lanka and the Maldives can be combined with recently published regional databases from the Pacific, Gulf, Atlantic, and Caribbean coasts of North America, Atlantic South America, Greenland, Antarctica, northwest Europe, the Barents Sea, the Mediterranean, China, Australia, New Zealand, other low-latitude locations, and high-resolution Common




Figure 1: Map showing the spatial distribution of sea-level data from different time periods: (A) the Last Glacial Maximum to present from new regional databases, (B) the Last Interglacial, (C) MIS 11, and (D) the Pliocene. References are given in Box 1.

Era reconstructions (Kopp et al. 2016; also Barnett et al. this issue; see Fig 1a). However, updates or further standardization may be required to fully integrate these recently published databases. Key spatial gaps remain in Arctic Canada, Pacific Central America, Pacific South America, and African coastlines, and there is a paucity of data spanning the deglacial period (i.e. older than 8 kyr).

The Last Interglacial

For the Last Interglacial, four primary databases collect Marine Isotope Stage (MIS) 5e sea-level indicators at global scale: 1) Kopp et al. (2009) included data from 42 locations and a variety of archives (isotopic, coral reef, geomorphological) and applied a standardized relationship to sea level at the time of formation (i.e. indicative meaning); 2) Pedoja et al. (2014) included data from 942 sites, however only elevation is reported (and often only mean elevations) without consideration of sample indicative meaning; 3) Dutton and Lambeck (2012) concentrated on coral reef archives from 16 sites (710 data points) and crucially standardized the U-series ages; 4) Hibbert et al. (2016) built on the Dutton and Lambeck dataset (32 locations, ca. 2,500 data points, for the last 700 kyr) adding additional standardization and coral depth distributions derived from modern ecological studies (Fig 1b). While at first glance there appears to be an abundance of Last Interglacial data, not all dated sea-level indicators have a full suite of database fields (for example, species dated, elevation, or reliable age determinations). Screening of the available databases suggests there are ca. 500 Last Interglacial sea-level indicators (excluding isotopic archives) with sufficient documentation to allow further analysis, 319 of which are located at 26 different locations on passive margins (Austermann et al. 2017). Databases from older time periods have often been “standalone” efforts with differing objectives, a major drawback of which is the varying way that archives have been interpreted with respect to past sea levels. One way forward is the approach taken by Rovere et al. (2016), where former sea levels are interpreted in terms of the entire geological or sedimentary facies (with ages derived from samples collected from within that facies), rather than considering each individually dated indicator separately.

Plio-Pleistocene interglacials

Beyond the Last Interglacial, there have been few attempts to compile and standardize sea-level data. Most compilations were completed to support modeling studies that did not focus on the creation of a database per se, and hence standardization is sometimes less rigorous than for Holocene and Last Interglacial proxies. For example, Creveling et al. (2017) report 38 sites dating to MIS 5a and MIS 5c to compare their elevation with glacial isostatic adjustment models. No attempt is made, however, at assessing or standardizing the indicative meaning of each proxy. For older interglacials, Bowen (2010) reported seven sites where MIS 11 shorelines

have been preserved (Fig 1c), and Rovere et al. (2015, 2014) estimated the indicative meaning for mid-Pliocene shorelines on the Atlantic coasts of the United States, South Africa, and South Australia (Fig 1d).

Future directions

Progress in improving the standardization of sea-level databases has also been accompanied by advancements in statistical and analytical methods used to infer spatial patterns and rates of RSL change from geological data that have a spatially and temporally sparse distribution and geochronological and elevational uncertainties (e.g. Austermann et al. 2017; Kopp et al. 2009, 2016). Future areas of development include more comprehensive and accurate use of data (e.g. incorporating non-Gaussian data distributions; see Hibbert et al. 2016), integration with physical models and their uncertainties (Milne et al. this issue) using machine learning approaches, and scaling spatio-temporal models to large geological datasets (Ashe et al. 2019).

Challenges remain in integrating databases compiled by different research groups over different time periods, and in developing cyberinfrastructure and open access visualization platforms to improve the longevity and accessibility of databases (e.g. Düsterhus et al. 2016). Improved understanding of the mechanisms driving RSL variability will be achieved through the standardization of sea-level databases, which will enhance the comparability and accessibility of information to improve both physical models and statistical reconstructions.

AFFILIATIONS

¹Asian School of the Environment, Nanyang Technological University, Singapore

²Research School of Earth Sciences, Australian National University, Canberra, Australia

³MARUM, University of Bremen, Germany

CONTACT

Nicole Khan: nicolekhan@ntu.edu.sg

REFERENCES

- Ashe EL et al. (2019) *Quat Sci Rev* 204: 58–77
 Austermann J et al. (2017) *Sci Adv* 3: e1700457
 Bowen DQ (2010) *Clim Past* 6: 19–29
 Creveling JR et al. (2017) *Quat Sci Rev* 163: 193–208
 Düsterhus A et al. (2016) *Clim Past* 12: 911–921
 Dutton A, Lambeck K (2012) *Science* 337: 216–219
 Hibbert FD et al. (2016) *Quat Sci Rev* 145: 1–56
 Hijma M et al. (2015) In: Shennan I et al. (Eds) *Handbook for Sea Level Research*. Wiley, 536–553
 Kopp RE et al. (2009) *Nature* 462: 863–867
 Kopp RE et al. (2016) *Proc Natl Acad Sci* 113: E1434–E1441
 Pedoja K et al. (2014) *Earth-Sci Rev* 132: 13–38
 Rovere A et al. (2014) *Earth Planet Sci Lett* 387: 27–33
 Rovere A et al. (2015) *Earth-Sci Rev* 145: 117–131
 Rovere A et al. (2016) *Earth-Sci Rev* 159: 404–427
 Shennan I, Horton B (2002) *J Quat Sci* 17: 511–526

Last Glacial Maximum to present	
Long et al. (2011)	Greenland
Vacchi et al. (2018a)	Atlantic Canada
Baranskaya et al. (2018)	Russian Arctic
Auriac et al. (2016)	Barents Sea
Lambeck et al. (2010)	Scandinavia
Briggs and Tarasov (2013)	Antarctica
Shennan et al. (2018)	British Isles
Hijma and Cohen (2019) Vink et al. (2007)	northwest Europe
García-Artola et al. (2018)	Atlantic Europe
Vacchi et al. (2014, 2016, 2018b) Shaw et al. (2018) Dean et al. (2019)	Mediterranean
Engelhart and Horton (2012) Hawkes et al. (2016)	US Atlantic
Engelhart et al. (2015) Reynolds and Simms (2015)	US Pacific
Hijma et al. (2015) Love et al. (2016)	Gulf of Mexico
Khan et al. (2017) Milne and Peros (2013)	Circum-Caribbean
Milne et al. (2005)	Atlantic South America
Cooper et al. (2019)	South Africa
Zong (2004)	China
Mann et al. (in review) Tam et al. (2018)	Southeast Asia, India, Sri Lanka, and the Maldives
Clement et al. (2016)	New Zealand
Lewis et al. (2013)	Australia
Hibbert et al. (2016, 2018)	Mid to low latitude locations
Khan et al. (2015)	Global
Last Interglacial	
Kopp et al. (2009)	Global
Pedoja et al. (2011, 2014)	Global
Hibbert et al. (2016)	Global
MIS 11	
Bowen et al. (2010)	see Figure 1
Pliocene	
Rovere et al. (2014, 2015)	see Figure 1

Box 1: Currently available standardized RSL databases. All data are shown in Figure 1. See online version of this article for links to full references (doi.org/10.22498/pages.27.1.10).