

## Full References

### PAGES Newsletter Vol.11, No 2 & 3, 2003:

---

There are references for the following articles:

- [Böhm](#)
  - [Ojala et al. Harvey](#)
  - [Goslar](#)
  - [Eronen et al.](#)
  - [Charman and Mäkilä](#)
  - [Bradley](#)
  - [Pilcher et al.](#)
  - [Widmann and Tett](#)
  - [Renssen and Osborne](#)
- 

Böhm Auer, I., Böhm, R. and Schöner, W., 2001: Austrian long-term climate 1767-2000 – Multiple instrumental climate time series from Central Europe. *Österr. Beitr. Zu Meteorologie und Geophysik* 25, 147 pp (plus Data- and Metadata- CD)

Böhm, R., Auer, I., Brunetti, M., Maugeri, M., Nanni, T. and Schöner, W., 2001: Regional temperature variability in the European Alps 1760-1998 from homogenized instrumental time series. *International Journal of Climatology* 21: 1779-1801

Jones P.D., 1994: Hemispheric surface air temperature variations: A reanalysis and an update to 1993. *Journal of Climate* 7: 1794-1802

Hanssen-Bauer, I. and Nordli, PØ., 1998: Annual and seasonal temperature variations in Norway 1876-1997. *DNMI reports, 25/98*, 29 pp.

Moberg, A. and Alexandersson, H., 1997: Homogenization of Swedish temperature data. Part II: Homogenised gridded air temperature compared with a subset of global air temperature since 1861. *International Journal of Climatology* 17: 35-54.

Peterson, T.C., Easterling, D.R., Karl, T.R., Groisman, P., Auer, I., Böhm, R., Plummer, N., Nicholis, N., Torok, S., Vincent, L., Tuomenvirta, H., Salinger, J., Førland, E.J., Hanssen-Bauer, I., Alexandersson, H., Jones, P. and Parker, D., 1998: Homogeneity Adjustments of In Situ Climate Data: A Review. *International Journal of Climatology* 18: 1493-1517

Vose, R.S., Peterson, T.C., Schmoyer, R.L., Eischeid, J.K., Steurer, P.M., Heim, R.R. and Karl, T.R., 1993: The Global Historical Climatology Network: Long-term monthly temperature, precipitation, and pressure data. Fourth AMS Symposium on Global Change Studies, Anaheim, CA., January 17-22 1993.

.Ojala et al. Hughen, K.A., Overpeck, J.T., Overpeck, J.T. and Anderson, R.F., 2000: Recent warming in a 500-year palaeotemperature record from varved sediments,

Upper Soper Lake, Baffin Island, Canada. *The Holocene* 10: 9-19.

Hurrell, J.W., 1995: Decadal trends in the North Atlantic Oscillation: Regional temperatures and precipitation. *Science* 291: 603-604.

Kuusisto, E., 1984: Snow accumulation and snowmelt in Finland. *Publications of the Water Research Institute* 55, 142 pp.

Luterbacher, J., Xoplaki, E., Dietrich, D., Jones, P.D., Davies, T.D., Portis, D., Gonzalez-Rouco, J.F., von Storch, H., Gyalistras, D., Casty, C., Wanner, H., 2002: Extending North Atlantic Oscillation reconstructions back to 1500. *Atmospheric Science Letters* 2: 114-124.

Ojala, A.E.K., Saarinen, T. and Salonen, V.-P., 2000: Preconditions for the formation of annually laminated lake sediments in southern and central Finland. *Boreal Environment Research* 5: 243-255.

Ojala, A.E.K. and Saarinen, T., 2002: Palaeosecular variation of Earth's magnetic field during the last 10,000 yrs based on the annually laminated sediment of Lake Nautajärvi, central Finland. *The Holocene* 12: 391-400.

Ojala, A.E.K. and Francus, P., 2002: X-ray densitometry vs. BSE -image analysis of thin-sections: a comparative study of varved sediments of Lake Nautajärvi, Finland. *Boreas* 31: 57-64.

Ojala, A.E.K., Heinsalu, A., Saarnisto, M., Tiljander, M., Annually laminated sediments date the drainage of the Ancylus Lake and early Holocene shoreline displacement in central Finland. *Quaternary International*, in press

Ojala, A.E.K. and Tiljander, M., Testing the fidelity of sediment chronology: comparing varve and paleomagnetic results from Holocene lake sediments from central Finland. *Quaternary Science Reviews*, in press

Petterson, G., 1996: Varved sediments in Sweden: a brief review. In Kemp, A. E. S (ed.) *Palaeoclimatology and Palaeoceanography from Laminated sediments*. Geol. Soc. London, Spec. Publication No.116. pp.73-77.

Petterson, G., Renberg, I., Anderson, N.J. and Segerström, U., 1999: A 6400 year annual record of minerogenic matter accumulation in Lake Kassjön, N. Sweden (a manuscript). In: Petterson, G. *Image analysis, varved lake sediments and climate reconstructions*. Ph.D-thesis, Department of Ecology and Environmental Science, Umeå University.

Renberg, I., 1982: Varved lake sediments - geochronological records of the Holocene. *Geologiska Föreningens i Stockholm Förhandlingar* 104: 275-279.

Saarinen, T. and Petterson, G., 2001: Image analysis techniques. In Last & Smol (eds) *Tracking Environmental Change Using Lake Sediments: Volume 2: Physical and*

Geochemical Methods, Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 23-39.

Snowball, I., Sandgren, P. and Petterson, G., 1999: The mineral magnetic properties of an annually laminated Holocene lake-sediment sequence in northern Sweden. *The Holocene* 9: 353–362.

Snowball, I.F. and Sandgren, P., 2002: Geomagnetic field variations in northern Sweden during the Holocene quantified from varved lake sediments and their implications for cosmogenic nuclide production rates. *The Holocene* 12: 517–530.

Tiljander, M., Ojala, A., Saarinen, T. and Snowball, I., 2002: Documentation of the physical properties of annually laminated (varved) sediments at a sub-annual to decadal resolution for environmental interpretation. *Quaternary International* 88: 5-12.

Turner, G.M. and Thompson, R. 1981: Lake sediment record of the geomagnetic secular variation in Britain during Holocene times. *Geophysical Journal of the Royal Astronomical Society* 65: 703-725.

Vehviläinen, B. and Lohvansuu, J., 1991: The effects of climate change on discharges and snow cover in Finland. *Hydrological Sciences - Journal - des Sciences Hydrologiques* 36: 109-121.

Vehviläinen, B. and Huttunen, M., 1997: Climate change and water resources in Finland. *Boreal Environment Research* 2: 3-18.

Zillén, L.M, Wastegård, S. and Snowball, I.F., 2002: Calendar year ages of three mid-Holocene tephra layers identified in varved lake sediments in west central Sweden. *Quaternary Science Reviews* 21: 1583–1591.

Zillén, L., Snowball, I.F., Sandgren, P. and Stanton, T., Occurrence of varved lake sediment sequences in Värmland, west central Sweden: lake characteristics, varve chronology and AMS radiocarbon dating. *Boreas*, in press (scheduled 3/2003).

Goslar, E., Raisbeck, G.M., Yiou, F. and Jouzel, J., 1997: Solar modulation of cosmogenic nuclide production over the last millennium: comparison between  $^{14}\text{C}$  and  $^{10}\text{Be}$  records. *Earth and Planetary Science Letters* 150: 453-462.

Goslar, T., 2001: Absolute production of radiocarbon and the long-term trend of atmospheric radiocarbon. *Radiocarbon* 43: 743-749.

Lal, D., 1988: Theoretically expected variations in the terrestrial cosmic-ray production of isotopes. [w:] *Solar-Terrestrial relationships and the Earth environment in the last millenia*. *Societa Italiana di Fisica* 95: 216-233.

Stuiver, M. and Braziunas, T., 1988: The solar component of the atmospheric  $^{14}\text{C}$  record. In: Stephenson, F.R. and Wolfendale, A.W. (eds): *Secular Solar and*

Geomagnetic Variations in the Last 10,000 Years, Kluwer Academic Publishers, 245-266.

Stuiver, M. and Braziunas, T., 1993: Sun, ocean, climate and atmospheric  $^{14}\text{C}$ : an evaluation of causal and spectral relationships. *The Holocene* 3: 289-305.

Stuiver, M. and Quay, P.D., 1980: Changes in Atmospheric Carbon-14 Attributed to a Variable Sun. *Science* 207: 11-19.

Stuiver, M., Braziunas, T., Becker, B. and Kromer, B., 1991: Climatic, solar, oceanic, and geomagnetic influences on Late-Glacial and Holocene atmospheric  $^{14}\text{C}/^{12}\text{C}$  change. *Quaternary Research* 35: 1-24.

Stuiver, M., Reimer, P.J., Bard, E., Warren Beck, J., Burr, G.S., Hughen, K.A., Kromer, B., McCormac, G., vanderPlicht, J. and Spurk, M., 1998: INTCAL98 Radiocarbon age calibration, 24,000-0 cal BP. *Radiocarbon* 40: 1041-1083.

Eronen et al. Eronen, M., Zetterberg, P., Briffa, K., Lindholm, M., Meriläinen, J., and Timonen, M., 2002: The supra-long Scots pine tree-ring record for northern Finnish Lapland – Part 1: chronology construction and initial inferences. *The Holocene* 12: 673-680.

Helama, S., Lindholm, M., Timonen, M. and Eronen, M., 2002: The supra-long Scots pine tree-ring record for northern Finnish Lapland – Part 2: interannual to centennial variability in summer temperatures for 7500 years. *The Holocene* 12: 681-687.

Lindholm, M., 1996: Reconstruction of the past climate from ring-width chronologies of Scots pine (*Pinus sylvestris* L.) at the northern forest limit in Fennoscandia. *Publications in Sciences* 40. PhD thesis. University of Joensuu. 169 p.

Lindholm, M., Eggertsson, Ó., Lovelius, N., Raspopov, O., Shumilov, O. and Läänelaid, A., 2001: Growth indices of North European Scots pine record the seasonal North Atlantic Oscillation. *Boreal Environment Research* 6: 275-284.

Lindholm, M., Kolström, T., Timonen, M. and Hallikainen, V., Growth variability and growth responses of forest-limit *Pinus sylvestris* in three age-classes in northern Fennoscandia. *Silva Fennica* (in press).

Lindholm, M., Lehtonen, H., Kolström, T., Meriläinen, J., Eronen, M. and Timonen, M., 2000: Climatic signals extracted from ring-width chronologies of Scots pine from the Northern, Middle and Southern parts of the boreal forest belt in Finland. *Silva Fennica* 34: 317-329.

Ogurtsov, M.G., Kocharov, G.E., Lindholm, M., Meriläinen, J., Eronen, M. and Nagovitsyn, Yu. A., 2002: Evidence of solar variation in tree-ring-based climate reconstructions. *Solar Physics* 205: 403-417.

.Charman and Mäkilä Barber, K.E., Chambers, F.M., Maddy, D., Stoneman, R. and

Brew, J.S., 1994: A sensitive high resolution record of late Holocene climatic change from a raised bog in northern England. *The Holocene* 4: 198-205.

Blackford, J.J. and Chambers, F.M., 1993: Determining the degree of peat decomposition in peat-based palaeoclimatic studies. *International Peat Journal* 5: 7-24.

Booth, R.K., 2002: Testate amoebae as paleoindicators of surface-moisture changes on Michigan peatlands: modern ecology and hydrological calibration. *Journal of Paleolimnology* 28: 329-348

Brenninkmeijer, C.A.M., Vangeel, B. and Mook, W.G., 1982: Variations in the d/h and o-18/o-16 ratios in cellulose extracted from a peat bog core. *Earth and Planetary Science Letters* 61: 283-290.

Campbell, I.D., Campbell, C., Yu, Z.C., Vitt, D.H. and Apps, M.J., 2000: Millennial-scale rhythms in peatlands in the western interior of Canada and in the global carbon cycle. *Quaternary Research* 54: 155-158.

Chambers, F.M. and Blackford, J.J., 2001: Mid- and late-Holocene climatic changes: a test of periodicity and solar forcing in proxy-climate data from blanket peat bogs. *Journal of Quaternary Science* 16: 329-338.

Chambers, F.M., Barber, K.E., Maddy, D. and Brew, J., 1997: A 5500-year proxy-climate and vegetation record from blanket mire at Talla Moss, Borders, Scotland. *The Holocene* 7: 391-399.

Charman, D.J., Brown, A.D., Hendon, D., Kimmel, A. and Karofeld, E., Testing the relationship between Holocene peatland palaeoclimate reconstructions and instrumental data. *Quaternary Science Reviews* (submitted).

Charman, D.J. and Hendon, D., 2000: Long-term changes in soil water tables over the past 4500 years: Relationships with climate and North Atlantic atmospheric circulation and sea surface temperature. *Climatic Change* 47: 45-59.

Charman, D.J., Caseldine, C., Baker, A., Gearey, B.R. and Hatton, J., 2001: Palaeohydrological records from peat profiles and speleothems in Sutherland, northwest Scotland. *Quaternary Research* 55: 223-234.

Gajewski, K., Vance, R., Sawada, M., Fung, I., Gignac, L.D., Halsey, L., John, J., Maisongrande, P., Mandell, P., Mudie, P.J., Richard, P.J.H., Sherin, R.A.G., Soroko, J. and Vitt, D., 2000: The climate of North America and adjacent ocean waters ca. 6 ka. *Canadian Journal of Earth Sciences* 37: 661-681.

Hendon, D., Charman, D.J. and Kent, M., 2001: Palaeohydrological records derived from testate amoebae analysis from peatlands in northern England: within-site variability, between-site comparability and palaeoclimatic implications. *The Holocene* 11: 127-148.

Hughes, P.D.M., Mauquoy, D., Barber, K.E. and Langdon, P.G., 2000: Mire-development pathways and palaeoclimatic records from a full Holocene peat archive at Walton Moss, Cumbria, England. *The Holocene* 10: 465-479.

Hughes, P.D.M., Mauquoy, D., Barber, K.E. and Langdon, P.G., 2000: Mire-development pathways and palaeoclimatic records from a full Holocene peat archive at Walton Moss, Cumbria, England. *The Holocene* 10: 465-479.

Langdon, P.G. and Barber, K.E., 2001: New Holocene tephtras and a proxy climate record from a blanket mire in northern Skye, Scotland. *Journal of Quaternary Science* 16: 753-759.

Mäkilä, M., 2001: Climate in relation to carbon accumulation on Kilpisuo, a raised bog in southern Finland. In: Vasiliev, S. V., Titlyanova, A. A., and Velichko, A. A. (eds.) *West Siberian peatlands and carbon cycle: Past and present: Proceedings of an International Symposium Noyabrsk*. pp.38-40.

Mäkilä, M., Saarnisto, M. and Kankainen, T., 2001: Aapa mires as a carbon sink and source during the Holocene. *Journal of Ecology* 89: 589-599.

Martinez-Cortizas, A., Pontevedra-Pombal, X., Garcia-Rodeja, E., Novoa-Munoz, J.C. and Shotyk, W., 1999: Mercury in a Spanish peat bog: Archive of climate change and atmospheric metal deposition. *Science* 284: 939-942.

Mauquoy, D., van Geel, B., Blaauw, M. and van der Plicht, J., 2002: Evidence from northwest European bogs shows 'Little Ice Age' climatic changes driven by variations in solar activity. *The Holocene* 12: 1-6.

McGlone, M.S. and Wilmshurst, J.M., 1999: A Holocene record of climate, vegetation change and peat bog development, east Otago, South Island, New Zealand. *Journal of Quaternary Science* 14: 239-254.

Mitchell, E.A.D., Van der Knaap, W.O., van Leeuwen, J.F.N., Buttler, A., Warner, B.G. and Gobat, J.M., 2001: The palaeoecological history of the Praz-Rodet bog (Swiss Jura) based on pollen, plant macrofossils and testate amoebae (Protozoa). *The Holocene* 11: 65-80.

Pendall, E., Markgraf, V., White, J.W.C., Dreier, M., and Kenny, R., 2001: Multiproxy record of late Pleistocene-Holocene climate and vegetation changes from a peat bog in Patagonia. *Quaternary Research* 55: 168-178.

Van Den Bogaard, C. and Schmincke, H.U., 2002: Linking the North Atlantic to central Europe: a high-resolution Holocene tephrochronological record from northern Germany. *Journal of Quaternary Science* 17: 3-20.

Wilmshurst, J.M., McGlone, M.S., and Charman, D.J., 2002: Holocene vegetation and climate change in southern New Zealand: Linkages between forest composition and quantitative surface moisture reconstructions from an ombrogenous bog. *Journal*

of Quaternary Science 17: 653-666.

Woodland, W.A., Charman, D.J. and Sims, P.C., 1998: Quantitative estimates of water tables and soil moisture in Holocene peatlands from testate amoebae. *The Holocene* 8: 261-273.

Xie, S., Nott, C.J., Avsejs, L.A., Volders, F., Maddy, D., Chambers, F.M., Gledhill, A., Carter, J.F. and Evershed, R.P., 2000: Palaeoclimate records in compound-specific delta D values of a lipid biomarker in ombrotrophic peat. *Organic Geochemistry* 31: 1053-1057.

Bradley, R.B., Mayewski, P.A., Sowers, T., Stuiver, M., Taylor, K.C. and Clark, P.U., 1997: Holocene climatic instability: a prominent, widespread event 8200 yr ago. *Geology* 25: 483-486.

Barber, D.C., Dyke, A., Hillaire-Marcel, C., Jennings, A.E., Andrews, J.T., Kerwin, M.W., Bilodeau, G., McNeely, R., Southon, J., Morehead, M.D., and Gagnon, J.M., 1999: Forcing of the cold event of 8,200 years ago by catastrophic drainage of Laurentide lakes. *Nature* 400: 344-348.

Bradley, R.S., 2003: Climate forcing during the Holocene. In: *Global Change in the Holocene: approaches to reconstructing fine-resolution climate change*. A.W. Mackay, R.W. Battarbee, H.J.B. Birks & F. Oldfield (eds.). Arnold, London (in press).

Crowley, T.J., 2000: Causes of climate change of the last 1000 years. *Science* 289: 270-277.

Rind, D., 2002: The Sun's role in climate variations. *Science* 296: 673-677.

Robock, A., 2000: Volcanic eruptions and climate. *Reviews of Geophysics* 38: 191-219.

Ruzmaikin, A.A., 1999: Can El Niño amplify the solar forcing of climate? *Geophysical Research Letters* 26: 2255-2258.

Zielinski, G., Mayewski, P.A., Meeker, L.D., Whitlow, S., Twickler, M.S., Morrison, M., Meese, D.A., Gow, A.J., and Alley, R.B., 1994: Record of explosive volcanism since 7000 B.C. from the GISP2 Greenland ice core and implications for the volcano-climate system. *Science* 264: 948-952. (See also comments and discussion, 1995: *Science* 267, 256-258).

Pilcher et al., van den Bogaard, P. and Schmincke, H.-U., 2002: Linking the North Atlantic to central Europe: a high resolution Holocene tephrochronological record from northern Germany. *Journal of Quaternary Science* 17: 3-20.

Dugmore, A.J., Larsen, G. and Newton, A.J., 1995: Seven isochrones in Scotland. *The Holocene* 5: 257-266.

- Haflidason, H., Eiriksson, J. and van Kreveland, S., 2000: The tephrochronology of Iceland and the North Atlantic region during the Middle and Late Quaternary: a review. *Journal of Quaternary Science* 15: 3-22.
- Hall, V.A. and Pilcher, J.R., 2002: Late quaternary Icelandic tephtras in Ireland and Great Britain: detection, characterization and usefulness. *The Holocene* 12: 223-230.
- Larsen, G. and Thorarinsson, S., 1977: H4 and other acid tephra layers. *Jokull* 27: 29-47.
- Pilcher, J.R. and Hall, V.A., 1992: Towards a tephrochronology for the Holocene of the north of Ireland. *The Holocene* 2: 255-259.
- Pilcher, J.R., Hall, V.A. and McCormac, F.G., 1996: An outline tephrochronology for the north of Ireland. *Journal of Quaternary Science* 11: 485-494.
- .Widmann and TettBarnett, T. B., 1999: Comparison of near-surface air temperature variability in 11 coupled global climate models. *Journal of Climate* 12: 511-518.
- Bauer, E., Claussen, M. and Brovkin, V., Assessing climate forcings of the Earth system for the past millennium. *Geophysical Research Letters*, accepted.
- Bertrand, C., Loutre, C.F., Crucifix, M. and Berger, A., 2002: Climate of the last millennium: a sensitivity study. *Tellus* 54A: 221-244.
- Collins, M., Tett, S.F.B. and Cooper, C., 2001: The internal climate variability of HadCM3, a version of the Hadley Centre coupled model without flux adjustments. *Climate Dynamics* 17: 61-81.
- Cubasch, U., Hegerl, G.C., Voss, R., Waszkewitz, J. and Crowley, T., 1997: Simulation with an O-AGCM of the influence of variations of the solar constant on the global climate. *Climate Dynamics* 13: 757-767.
- Crowley, T.J., 2000: Causes of climate change over the past 1000 years. *Science* 289: 270-277.
- Fischer-Bruns, I., Cubasch, U., von Storch, H., Zorita, E., González-Rouco, F. and Luterbacher, J., 2002: Modelling the Late Maunder Minimum with a 3-dimensional OAGCM. *CLIVAR Exchanges*, 7: 59-61. International CLIVAR Project Office, Southampton, UK, ISSN No. 1026-0471.
- Manabe, S. and Stouffer, R.J., 1996: Low-frequency variability of air temperature in a 1000-year integration of a coupled atmosphere-ocean-land surface model. *Journal of Climate* 9: 376-393.
- Hall, A. and Stouffer, R. J., 2001: An Extreme Climate Event in a Coupled Ocean-Atmosphere Simulation without External Forcing. *Nature* 409: 171-174.



Hegerl, G.C., von Storch, H., Hasselmann, K., Santer, B.D., Cubasch, U. and Jones, P.D., 1996: Detecting anthropogenic climate change with an optimal fingerprint method. *Journal of Climate* 9: 2281-2306.

Shindell, D.T., Schmidt, G.A. Mann, M.E. Rind, D. and Waple, A., 2001: Solar forcing of regional climate change during the Maunder Minimum. *Science* 294: 2149-2152.

Stott, P.A., Tett, S.F.B., Jones, G.S., Allen, M.R., Mitchell, J.F.B. and Jenkins, G. J., 2000: External control of 20th century temperature variations by natural and anthropogenic forcings. *Science* 15: 2133-2137.

Zorita, E., González-Rouco, F. and Legutke, S., 2003: Testing the Mann et al. (1998) approach to paleoclimatic reconstructions in the context of a 1000 year control simulation with the Echo-G coupled climate model. *Journal of Climate*, accepted.

Zorita, E. and González-Rouco, F., 2002: Are temperature proxies adequate for North Atlantic oscillation reconstructions? *Geophysical Research Letters* 29, 48-1 - 48-4.

.Renssen and OsborneBrovkin, V., Bendtsen, J., Claussen, M., Ganopolski, A., Kubatzki, C., Petoukhov, V. and Andreev, A., 2002: Carbon cycle, vegetation and climate dynamics in the Holocene: experiments with the CLIMBER-2 model. *Global Biogeochemical Cycles* 16: doi:10.1029/2001GB001662.

Collins, M., Osborn, T.J., Tett, S.F.B., Briffa, K.R. and Schweingruber, F.H., 2002: A comparison of the variability of a climate model with palaeo-temperature estimates from a network of tree-ring densities. *Journal of Climate* 15: 1497-1515.

Crowley, T.J., 2000: Causes of climate change over the past 1000 years, *Science* 289: 270-277.

Crucifix, M., Loutre, M.F., Tulkens, P., Fichefet, T. and Berger, A., 2002: Climate evolution during the Holocene: a study with an Earth system model of intermediate complexity. *Climate Dynamics* 19: 43-60.

Jones, J.M. and Widmann, M., 2002: Reconstructing large-scale variability from palaeoclimatic evidence by means of Data Assimilation Through Upscaling and Nudging (DATUN). KIHZ book, Springer, accepted.

Reichert, B.K., Bengtsson, L. and Oerlemans, J., 2001: Midlatitude forcing mechanisms for glacier mass balance investigated using general circulation models. *Journal of Climate* 14: 3767-3784.

Renssen, H., Goosse, H. and Fichefet, T., 2002: Modeling the effect of freshwater pulses on the early Holocene climate: the influence of high frequency climate variability. *Paleoceanography* 17, No. 2, 10.1029/2001PA000649.

Weber, S.L. and Oerlemans, J., 2002: Holocene glacier variability: three case studies using an intermediate-complexity climate model. *Holocene*, in press