

UCL Press Release

Under embargo until 12 October 2018, 10:00 London time (BST) / 5:00am US Eastern time / 18:00 Japanese time / 20:00 Australian Eastern Time

Does climate vary more from century to century when it is warmer?

Century-scale climate variability was enhanced when the Earth was warmer during the Last Interglacial period (129-116 thousand years ago) compared to the current interglacial (the last 11,700 years), according to a new UCL-led study.

The findings, published today in *Nature Communications* and funded by the Natural Environment Research Council (NERC) and the Australian Research Council (ARC), reveal that the Last Interglacial period was punctuated by a series of century-scale arid events in southern Europe and cold water-mass expansions in the North Atlantic.

Assessing natural climate variability under relatively warm conditions is crucial to inform projections under future carbon emission scenarios. Professor Chronis Tzedakis (UCL Geography), study lead author, said: "The Last Interglacial is particularly relevant because it provides insights into climate processes during a period of excess warmth."

The Last Interglacial period contained an interval of intense Arctic warming, with surface air temperatures estimated at 3-11°C above pre-industrial, comparable to high-latitude warming scenarios for the end of this century.

Global sea-level during the Last Interglacial is estimated to have been ~6-9 m above present, with 0.6-3.5 m derived from melting of the Greenland Ice Sheet.

Previously, several North Atlantic and European records have detected century-scale changes in temperature and precipitation within the Last Interglacial, but there has been considerable uncertainty over the timing, extent and origin of these climate oscillations.

This new study by international researchers from twelve institutions used marine and terrestrial geological archives, coupled with climate model experiments, to create the most detailed timeline of ocean and atmosphere changes in the North Atlantic and southern Europe during the Last Interglacial.

To address the uncertainties in comparing records from different environments, researchers produced a "stratigraphic 'rosetta stone' by analysing different fossils from the same

sediment samples in a marine core off Lisbon,” said Dr Luke Skinner (Cambridge University) who led the palaeoceanographic analyses.

“The marine core also contained pollen transported from the Tagus river into the deep sea, thus enabling a direct comparison of vegetation and North Atlantic ocean changes,” said Dr Vasiliki Margari (UCL Geography), who undertook the pollen analysis.

Changes in vegetation, primarily caused by variations in the amount of rainfall, were then linked to changes in the chemical signature of rainfall recorded in stalagmites from Corchia Cave in northern Italy.

“The Corchia record is particularly important because it is supported by very detailed radiometric dating using the decay of uranium isotopes, producing one of the best chronologies for this period available,” said Dr Russell Drysdale (University of Melbourne), who led the team studying the Italian cave.

Climate model experiments, undertaken by Dr Laurie Menviel and Dr Andrea Taschetto of the University of New South Wales Sydney, revealed that the spatial fingerprint of these changes was consistent with disruptions of the Atlantic meridional overturning circulation.

Greenland ice-melt and runoff as a result of strong high-latitude warming during the Last Interglacial may have contributed to the weakening of the Atlantic meridional overturning circulation and to the observed climate changes.

“Although not a strict analogue for future anthropogenically-driven changes, the profile of the Last Interglacial that emerges is one of enhanced century-scale climate instability, with implications for ice-sheet and ocean dynamics,” said Professor Tzedakis. “Future research efforts should focus on constraining further the extent of melting and runoff from the Greenland ice-sheet and its effects on ocean circulation during the Last Interglacial.”

Notes to Editors

For more information or to speak to the researchers involved, please contact Dr Rebecca Caygill, UCL Media Relations. T: +44 (0)20 3108 3846 / +44 (0)7733 307 596, E: r.caygill@ucl.ac.uk

‘Enhanced climate instability in the North Atlantic and southern Europe during the Last Interglacial’ by P.C. Tzedakis et al. will be published in *Nature Communications* on Friday 12

October 2018, 10:00 London time (BST) / 5:00am US Eastern time/18:00 Japanese time / 20:00 Australian Eastern Time and is under a strict embargo until this time.

The DOI for this paper will be 10.1038/s41467-018-06683-3

About UCL (University College London)

UCL was founded in 1826. We were the first English university established after Oxford and Cambridge, the first to open up university education to those previously excluded from it, and the first to provide systematic teaching of law, architecture and medicine. We are among the world's top universities, as reflected by performance in a range of international rankings and tables. UCL currently has over 41,000 students from 150 countries and over 13,500 staff. Our annual income is more than £1 billion.

www.ucl.ac.uk | Follow us on Twitter [@uclnews](https://twitter.com/uclnews) | Watch our YouTube channel [YouTube.com/UCLTV](https://www.youtube.com/UCLTV)